







# Numerical integration

Calculation of definite integrals implies to replace the function by another one (usually a polynomial form) which is a good approximation and easier to compute.  **Error estimation** depends on parameters of the method and refinement of spatial grid discretization. It is interesting to check these errors using various algorithm and mesh resolution.

- Equally spaced methods :
  -  [Numerical\\_integration](#)
  -  [Trapezoidal\\_rule](#)
  -  [Newton-Cotes\\_formulas](#)
  -  [Simpson's rule and composite Simpson's rule](#)
- If intervals between interpolation points vary :
  -  [Gaussian\\_quadrature](#)

## Références

- Numerical recipes, The Art of Scientific Computing 3rd Edition, William H. Press, Saul A. Teukolsky, William T. Vetterling, Brian P. Flannery, 2007, isbn: 9780521880688
  - <http://numerical.recipes/>
  - [http://www2.units.it/ipi/students\\_area/imm2/files/Numerical\\_Recipes.pdf](http://www2.units.it/ipi/students_area/imm2/files/Numerical_Recipes.pdf), p 129...
  - <http://apps.nrbook.com/empanel/index.html#>

From:  
<https://dvillers.umons.ac.be/wiki/> - **Didier Villers, UMONS - wiki**

Permanent link:  
[https://dvillers.umons.ac.be/wiki/teaching:methcalchim:numerical\\_integration?rev=1480080910](https://dvillers.umons.ac.be/wiki/teaching:methcalchim:numerical_integration?rev=1480080910)

Last update: **2016/11/25 14:35**

